



Axsome Therapeutics Highlights Commitment to Innovating Treatments for Mental Health Conditions with Presentations at Psych Congress 2023

September 7, 2023

Data presentations explore effects on patient-centric efficacy measures of treatments for patients with depression, and excessive sleepiness associated with obstructive sleep apnea and narcolepsy

NEW YORK, Sept. 07, 2023 (GLOBE NEWSWIRE) -- Axsome Therapeutics, Inc. (NASDAQ: AXSM), a biopharmaceutical company developing and delivering novel therapies for the management of central nervous system (CNS) disorders, today announced presentations of data from its CNS portfolio at the 2023 U.S. Psych Congress, being held in Nashville, Tenn. from September 6-10. The presentations demonstrate the Company's commitment to continued innovation to address unmet needs in difficult-to-treat CNS disorders.

"Axsome's presentations at Psych Congress highlight our commitment to advancing research of medicines that have the potential to improve the lives of patients with brain health conditions such as depression and sleep disorders," said Cecilia Brain, MD, PhD, Vice President of Medical Affairs at Axsome. "These presentations explore the meaningful impact that our treatments could have for patients and underscore the unique pharmacology of our novel medicines which may be relevant to multiple clinical conditions."

The presentations will include eight posters with data and analyses which examine the effects of AUVELITY® (dextromethorphan-bupropion) on sleep quality and cognitive and physical functioning in patients with major depressive disorder (MDD), and the effects of SUNOSI® (solriamfetol) on cognition in patients with excessive daytime sleepiness (EDS) due to obstructive sleep apnea (OSA). Five of the eight posters are finalists under consideration for the Psych Congress Poster Awards.

Key highlights include:

- Data examining durability of efficacy and tolerability with long-term (up to 12 months) treatment with AUVELITY in patients with MDD, including measures of depressive symptoms, anxiety, and cognitive and physical functioning.
- Data examining the effect of AUVELITY on depression-related disability, and patient-reported measures of insomnia in patients with MDD.
- Data examining treatment effect sizes of SUNOSI in patients with EDS associated with narcolepsy and OSA.
- Data examining the effects of SUNOSI on cognition in patients with EDS and OSA.

All posters will be presented on September 8 and 9 from 1:30-3 p.m. Central Time in Exhibit Hall A and B. An asterisk ahead of the title (*) indicates presentations under consideration as finalists for the Psych Congress Poster Awards, being presented from 6:45-8 p.m. Central Time on September 9. Details for the presentations are as follows:

Major Depressive Disorder:

*** Title:** AXS-05 (Dextromethorphan-Bupropion) Significantly Improved Functioning in Major Depressive Disorder: Analysis of the Domains of the Sheehan Disability Scale

Lead Author: Andrew Cutler, MD, SUNY Upstate Medical University and Neuroscience Education Institute

Poster Number: 44

*** Title:** Improvements in Cognitive and Physical Functioning Outcomes in Depressed Individuals Treated with AXS-05 (Dextromethorphan-Bupropion): Results from the EVOLVE Open-Label, Long-Term Study

Lead Author: Maurizio Fava, MD, Director, Division of Clinical Research, Massachusetts General Hospital (MGH) Research Institute

Poster Number: 51

Title: Assessment of Withdrawal Symptoms After Discontinuation of AXS-05 (Dextromethorphan-Bupropion) Treatment: Results From the GEMINI Trial

Lead Author: Rakesh Jain, MD, Texas Tech University School of Medicine-Permian Basin

Poster Number: 71

Title: Impact of AXS-05 (Dextromethorphan-Bupropion) on Patient-Reported Insomnia Symptoms: Results From the GEMINI Trial

Lead Author: Manish Jha, MD, O'Donnell Clinical Neuroscience Scholar Center for Depression Research and Clinical Care

Poster Number: 73

Excessive Daytime Sleepiness in Obstructive Sleep Apnea and Narcolepsy:

*** Title:** Effects of Solriamfetol on Cognitive Function in Participants With Cognitive Impairment Associated With Excessive Daytime Sleepiness in Obstructive Sleep Apnea: Results of the SHARP Study

Lead Author: Hans Van Dongen, PhD, Washington State University

Poster Number: 32

*** Title:** Solriamfetol Improves Cognitive Performance in Preclinical Models of Sleep Apnea and in a Randomized Placebo-Controlled Study of Sleep Apnea Participants (SHARP)

Lead Author: David Gozal, MD, University of Missouri

Poster Number: 33

*** Title:** Solriamfetol for Excessive Sleepiness in Narcolepsy and Obstructive Sleep Apnea: Effect Sizes and Numbers Needed to Treat or Harm

Lead Author: Craig Chepke, MD, SUNY Upstate Medical University

Poster Number: 45

Title: SURVEY Study of Solriamfetol: Initiation, Titration, Safety, Efficacy, and Follow-Up Experience for Patients With OSA in Germany

Lead Author: Yaroslav Winter, MD, Mainz Comprehensive Epilepsy and Sleep Medicine Center, Department of Neurology, Johannes Gutenberg-University, Mainz, Germany

Poster Number: 99

INDICATION AND IMPORTANT SAFETY INFORMATION

WHAT IS AUVELITY (aw-VEHL-ah-tee)? It is a prescription oral medicine used to treat adults with major depressive disorder (MDD). It is not known if Auvelity is safe and effective for use in children.

Auvelity is not approved for uses other than the treatment of MDD. The ingredients in Auvelity, bupropion and dextromethorphan, are the same ingredients found in some other medicines approved for other uses.

WHAT IS THE MOST IMPORTANT INFORMATION I SHOULD KNOW ABOUT AUVELITY?

Auvelity and other antidepressant medicines may increase suicidal thoughts and actions in some children, adolescents, and young adults, especially within the first few months of treatment or when the dose is changed. Auvelity is not for use in children.

You should pay close attention to any new or sudden changes in mood, behavior, thoughts, or feelings or if you develop suicidal thoughts or actions. This is very important when starting or changing the dose of an antidepressant medicine.

Call your healthcare provider (HCP) or get emergency help right away if you or your loved one have any of the following symptoms, especially if they are new, worse, or worry you:

- suicidal thoughts or actions
- new or worsening depression or anxiety
- agitation or restlessness
- trouble sleeping (insomnia)
- acting aggressive, being angry violent
- an extreme increase in activity and talking (mania)
- panic attacks
- new or worsening irritability
- acting on dangerous impulses
- other unusual changes in behavior or mood

Do not take Auvelity if you:

- have or had a seizure disorder.
- have or had an eating disorder like anorexia or bulimia.
- have recently and suddenly stopped drinking alcohol or use medicines called benzodiazepines, barbiturates, or anti-seizure medicines, and you have recently suddenly stopped taking them.
- are taking a monoamine oxidase inhibitor (MAOI), have stopped taking an MAOI in the last 14 days, or are being treated with the antibiotic linezolid or intravenous methylene blue. Ask your HCP or pharmacist if you are unsure whether you take an MAOI. Do not start taking an MAOI until you have stopped taking Auvelity for at least 14 days.
- are allergic to dextromethorphan, bupropion, or any other ingredients in Auvelity.

Auvelity may cause serious side effects. Ask your HCP how to recognize the serious side effects below and what to do if you think you have one:

Seizures. There is a risk of seizures during treatment with Auvelity. The risk is higher if you take higher doses of Auvelity, have certain medical problems, or take Auvelity with certain other medicines. Do not take Auvelity with other medicines unless your healthcare provider tells you to.

If you have a seizure during treatment with Auvelity, stop taking Auvelity and call your HCP right away. Do not take Auvelity again if you have a seizure. Increases in blood pressure (hypertension). Some people may get high blood pressure during treatment with Auvelity. Your HCP should check your blood pressure before you start taking and during treatment with Auvelity.

Manic episodes. Manic episodes may happen in people with bipolar disorder who take Auvelity. Symptoms may include:

- greatly increased energy
- racing thoughts
- unusually grand ideas

- talking more or faster than usual
- severe trouble sleeping
- reckless behavior
- excessive happiness or irritability

Unusual thoughts or behaviors. One of the ingredients in Auvelity (bupropion) can cause unusual thoughts or behaviors, including delusions (believing you are someone else), hallucinations (seeing or hearing things that are not there), paranoia (feeling that people are against you), or feeling confused. If this happens to you, call your HCP.

Eye problems (angle-closure glaucoma). Auvelity may cause a type of eye problem called angle-closure glaucoma in people with certain other eye conditions. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are. Call your HCP if you have eye pain, changes in your vision, or swelling or redness in or around the eye.

Dizziness. Auvelity may cause dizziness which may increase your risk for falls.

Serotonin syndrome. A potentially life-threatening problem called serotonin syndrome can happen when you take Auvelity with certain other medicines. Call your HCP or go to the nearest hospital emergency room right away if you have any of the following signs and symptoms:

- agitation
- hallucinations
- confusion
- coma
- fast heartbeat
- blood pressure changes
- dizziness
- sweating
- flushing
- high body temperature (hyperthermia)
- shaking (tremors), stiff muscles, or muscle twitching
- loss of coordination
- seizures
- nausea, vomiting, diarrhea

COMMON SIDE EFFECTS

The most common side effects of Auvelity include dizziness, headache, diarrhea, feeling sleepy, dry mouth, sexual function problems, and excessive sweating.

These are not all the possible side effects of Auvelity. Tell your doctor if you have any side effects. You can report side effects at 1-800-FDA-1088 or www.fda.gov/medwatch.

BEFORE USING

- Tell your HCP about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.
- It is important to tell your HCP if you are taking:
 - other medicines containing bupropion or dextromethorphan
 - medicines to treat depression, anxiety, psychotic or thought disorders, including selective serotonin reuptake inhibitors (SSRIs) and tricyclic antidepressants
 - theophylline
 - corticosteroids
 - oral diabetes medicines or use insulin to control your blood sugar
 - medicines to control appetite (anorectic)
 - nicotine medicines to help you stop smoking
 - street (illicit) drugs
 - benzodiazepines, sedative-hypnotic (sleep medicines), or opiates
- If you are unsure if you take any of these medicines, ask your HCP. They can tell you if it is safe to take Auvelity with your other medicines.
- Tell your HCP if you are pregnant or plan to become pregnant. Auvelity may harm your unborn baby if you take it during pregnancy. Auvelity is not recommended during pregnancy. Your HCP will prescribe another treatment for females who plan to become pregnant.

- One of the ingredients in Auvelity passes into your breast milk. Do not breastfeed during treatment with Auvelity and for 5 days after the final dose.

Tell your HCP about all your medical conditions, including if you:

- have problems with your liver or kidneys.
- have diabetes, heart disease, or high blood pressure.
- have a history of seizure, stroke, eating disorder, head injury, or have a tumor in your brain or spinal cord.
- have a history of alcohol or drug abuse.
- have a history of seizure, eating disorder, or abuse alcohol or drugs.
- have low blood sugar, low blood sodium levels, or a history of falls.
- you take certain other medicines that could interact with Auvelity.
- have or had a condition known as bipolar disorder, a family history of bipolar disorder, suicide, or depression.
- have high pressure in the eye (glaucoma).

Review the list below with your HCP. Auvelity may not be right for you if:

- you drink a lot of alcohol.
- you abuse prescription or street drugs.
- you are pregnant or plan to become pregnant.
- you are breastfeeding or plan to breastfeed.

HOW TO TAKE

- Auvelity is available by prescription only.
- Take Auvelity exactly as instructed by your HCP.
- Take Auvelity 1 time a day for 3 days, then increase your dose to 2 times a day (taken at least 8 hours apart). Do not take more than 2 Auvelity tablets in 24 hours.
- If you miss a dose, do not take an extra dose. Wait and take your next dose at the regular time. Do not take more than 1 dose of Auvelity at a time.
- Do not change your dose or stop taking Auvelity without talking to your HCP.
- Swallow Auvelity tablets whole. Do not crush, chew, or divide the tablets.
- Do not give Auvelity to other people.
- If you take too much Auvelity call your HCP or seek medical advice promptly.

LEARN MORE

For more information about Auvelity, call 866-496-2976 or visit [Auvelity.com](https://www.auvelity.com).

This summary provides basic information about Auvelity but does not include all information known about this medicine. Read the information that comes with your prescription each time your prescription is filled. This information does not take the place of talking with your doctor. Be sure to talk to your doctor or other HCP about Auvelity and how to take it. Your HCP is the best person to help you decide if Auvelity is right for you.

AUV CON BS 10/2022

Please see full [Prescribing Information](#), including Boxed Warning for suicidal thoughts and behaviors, and [Medication Guide](#).

More information about Sunosi, including Full Prescribing Information and Medication Guide, is available [here](#).

Important Safety Information

Before taking SUNOSI, tell your doctor about all of your medical conditions, including if you:

- have heart problems, high blood pressure, kidney problems, diabetes, or high cholesterol.
- have had a heart attack or a stroke.
- have a history of mental health problems (including psychosis and bipolar disorders), or of drug or alcohol abuse or addiction.
- are pregnant or planning to become pregnant. It is not known if SUNOSI will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if SUNOSI passes into your breast milk. Talk to your doctor about the best way to feed your baby if you take SUNOSI.

Do not take SUNOSI if you are taking, or have stopped taking within the past 14 days, a medicine used to treat depression called a monoamine oxidase inhibitor (MAOI).

What are the possible side effects of SUNOSI?

SUNOSI may cause serious side effects, including:

- **Increased blood pressure and heart rate.** SUNOSI can cause blood pressure and heart rate increases that can increase

the risk of heart attack, stroke, heart failure, and death. Your doctor should check your blood pressure before, and during, treatment with SUNOSI. Your doctor may decrease your dose or tell you to stop taking SUNOSI if you develop high blood pressure that does not go away during treatment with SUNOSI.

- **Mental (psychiatric) symptoms including anxiety, problems sleeping (insomnia), irritability, and agitation.** Tell your doctor if you develop any of these symptoms. Your doctor may change your dose or tell you to stop taking SUNOSI if you develop side effects during treatment with SUNOSI.

The most common side effects of SUNOSI include:

- headache
- decreased appetite
- problems sleeping
- nausea
- anxiety

These are not all the possible side effects of SUNOSI. Call your doctor for advice about side effects.

SUNOSI (solriamfetol) is available in 75 mg and 150 mg tablets and is a federally controlled substance (CIV) because it contains solriamfetol that can be a target for people who abuse prescription medicines or street drugs. Keep SUNOSI in a safe place to protect it from theft. Never give or sell your SUNOSI to anyone else because it may cause death or harm them and it is against the law. Tell your doctor if you have ever abused or been dependent on alcohol, prescription medicines, or street drugs.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see here for full [Prescribing Information](#)

SUN CON ISI 05/2022

About Axsome Therapeutics, Inc.

Axsome Therapeutics, Inc. is a biopharmaceutical company developing and delivering novel therapies for central nervous system (CNS) conditions that have limited treatment options. Through development of therapeutic options with novel mechanisms of action, we are transforming the approach to treating CNS conditions. At Axsome, we are committed to developing products that meaningfully improve the lives of patients and provide new therapeutic options for physicians. For more information, please visit the Company's website at axsome.com. The Company may occasionally disseminate material, nonpublic information on the company website.

Forward-Looking Statements

Certain matters discussed in this press release are "forward-looking statements". We may, in some cases, use terms such as "predicts," "believes," "potential," "continue," "estimates," "anticipates," "expects," "plans," "intends," "may," "could," "might," "will," "should" or other words that convey uncertainty of future events or outcomes to identify these forward-looking statements. In particular, the Company's statements regarding trends and potential future results are examples of such forward-looking statements. The forward-looking statements include risks and uncertainties, including, but not limited to, the continued commercial success of our Sunosi® and Auvelity® products and the success of our efforts to obtain any additional indication(s) with respect to solriamfetol and/or AXS-05; the success, timing and cost of our ongoing clinical trials and anticipated clinical trials for our current product candidates, including statements regarding the timing of initiation, pace of enrollment and completion of the trials (including our ability to fully fund our disclosed clinical trials, which assumes no material changes to our currently projected revenues or expenses), futility analyses and receipt of interim results, which are not necessarily indicative of the final results of our ongoing clinical trials, and/or data readouts, and the number or type of studies or nature of results necessary to support the filing of a new drug application ("NDA") for any of our current product candidates; our ability to fund additional clinical trials to continue the advancement of our product candidates; the timing of and our ability to obtain and maintain U.S. Food and Drug Administration ("FDA") or other regulatory authority approval of, or other action with respect to, our product candidates, including statements regarding the timing of any NDA submission; whether issues identified by FDA in the complete response letter may impact the potential approvability of the Company's NDA for AXS-07 for the acute treatment of migraine in adults with or without aura, pursuant to our special protocol assessment for the MOMENTUM clinical trial; the Company's ability to successfully defend its intellectual property or obtain the necessary licenses at a cost acceptable to the Company, if at all; the successful implementation of the Company's research and development programs and collaborations; the success of the Company's license agreements; the acceptance by the market of the Company's products and product candidates, if approved; the Company's anticipated capital requirements, including the amount of capital required for the continued commercialization of Sunosi and Auvelity and for the Company's commercial launch of its other product candidates, if approved, and the potential impact on the Company's anticipated cash runway; unforeseen circumstances or other disruptions to normal business operations arising from or related to COVID-19; and other factors, including general economic conditions and regulatory developments, not within the Company's control. The factors discussed herein could cause actual results and developments to be materially different from those expressed in or implied by such statements. The forward-looking statements are made only as of the date of this press release and the Company undertakes no obligation to publicly update such forward-looking statements to reflect subsequent events or circumstance.

Axsome Contacts:

Investors:
Mark Jacobson
Chief Operating Officer
Axsome Therapeutics, Inc.
One World Trade Center, 22nd Floor
New York, NY 10007
Tel: 212-332-3243

Email: mjacobson@axsome.com
www.axsome.com

Media:

Darren Opland
Director, Corporate Communications
Axsome Therapeutics, Inc.
One World Trade Center, 22nd Floor
New York, NY 10007
Tel: 929-837-1065
Email: dopland@axsome.com
www.axsome.com



Source: Axsome Therapeutics, Inc.